DR. JASON PIROZZOLO

A Sports Medicine View of Football Injuries: What Matters First?

A player drifts down a quiet New Town street before dinner, running a route while a football hangs against the fading light. One step lands over the crooked storm grate, the foot catches, and the whole play changes in a breath. The ball skips away, and the next few seconds tell more than the throw did.

The short answer is that most football injuries are minor, but the first visit is about sorting out what you can safely watch and what needs a closer exam. "I am looking for the story first, then what your body can and cannot do right now," says Jason Pirozzolo, DO, a sports medicine physician. You want a plain answer about what matters, what does not, and what the next few days should look like.

What points one way and what points the other

The first pass is usually simple. A twist over that crooked grate points Dr. Pirozzolo toward an ankle, knee, or thigh problem, while a direct blow points more toward a bruise in the exact place that took the hit.

Football also brings a few patterns that matter. A direct hit to the front of the thigh fits a quadriceps contusion, meaning a deep thigh bruise, and that can keep you out longer if it is not handled well. A blow over the top of the hip fits a hip pointer, meaning a painful bruise over the hip bone, and that often takes weeks to calm down. "The mechanism helps narrow it fast," Dr. Pirozzolo says. "How it happened usually tells me where to start."

Neck and shoulder symptoms have their own clues. A stinger, meaning a brief nerve stretch injury, often happens when the shoulder is forced down while the neck bends the other way. Concussion is also a common football concern, and Dr. Pirozzolo says a possible concussion needs to be recognized and checked with standard sideline tools instead of being brushed off as just getting your bell rung. "A helmet helps with some head and face injuries, but it does not protect the neck from every football hit," he says.

How long recovery usually lasts

Recovery can be short or it can stretch out, and the reason is usually the kind of injury rather than your toughness. Many football injuries are minor, but bruises in the thigh and hip can linger, and a concussion often needs a more careful pace with early mental and physical rest.

That means two players who look similar at first can have different timelines. A simple bruise may settle in days, while a hip pointer can take weeks, and a concussion recovery depends on how symptoms behave as normal thinking and movement return. "I tell people not to judge the next month by the first hour," Dr. Pirozzolo says. "The body usually declares itself after the exam and the first few days."

What changes the range is whether pain is local or spreading, whether strength comes back evenly, and whether normal movement returns without a limp or guarded motion. For a suspected concussion, early rest is part of care. For a knee that has already had ACL reconstruction, meaning surgery to rebuild a torn knee ligament, Dr. Pirozzolo is careful because functional gaps can still be present after a player has been cleared.

Getting back to football and daily life

Getting back starts with normal daily movement first, then football work later. Before you think about the next throw, cut, or tackle, Dr. Pirozzolo wants to see that you can walk, turn, and load the injured side without a protective hitch.

For shoulder problems, return testing may look at side to side strength, power, and how well you support yourself through the arms in a closed-chain test, meaning your hand stays planted while you move. For a knee after ACL recovery, one common benchmark from a 1991 study by Noyes and colleagues is that the involved leg should do at least 90 percent as well as the other leg on hop testing. "Cleared is not the same as ready," Dr. Pirozzolo says. "I want to see clean movement, confidence, and good side to side performance."

- Walk normally through the day before sport drills.
- Stop if pain makes you change your stride or motion.
- Rest early after a possible concussion, until Dr. Pirozzolo tells you how to progress.
- Do not judge a shoulder or knee by pain alone. Compare strength and control side to side.

That same New Town street scene is a good reminder. The trouble started with one awkward step, and the return should be just as specific, with the same motions tested before full football work resumes.

"What matters most is not the label people use on the sideline," Dr. Pirozzolo says. "What matters is whether the pattern fits, the exam makes sense, and your movement has come back." If that step over the grate still shows up in the way you walk or turn after the next day or two, have it checked. To be seen,

If that step over the grate still shows up in the way you walk or turn after the next day or two, have it checked. To be seen, contact Key West Concierge Orthopedics.

Questions About Football Injuries

Do most football injuries need a long recovery?

No. Many football injuries are minor, but some bruises, concussions, and ligament problems need a slower plan.

What does Dr. Pirozzolo care about most at the first visit?

He starts with how the injury happened, where the pain is, and what you can do right now without a limp, weakness, or guarded motion.

When are you ready to return to football?

You are closer when normal daily movement has returned and your strength, control, and confidence match the uninjured side well enough for sport testing.

Sources

- Shea K, Nilsson K. Youth Sports and Related Injuries. Textbook chapter, year not provided in excerpt.
- Youth Sports and Related Injuries. Textbook chapter, year not provided in excerpt.
- Patricios J, et al. Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport-Amsterdam, October 2022. Consensus statement, British Journal of Sports Medicine, 2023.
- Pediatric Orthopedics. Textbook chapter excerpt, year not provided in excerpt.
- Myer GD, et al. Utilization of modified NFL combine testing to identify functional deficits in athletes following ACL reconstruction. Primary study summarized in textbook chapter, 2011.
- Rotator Cuff Injuries. Textbook chapter excerpt, year not provided in excerpt.
- Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. Primary study summarized in textbook chapter, 1991.

Medically reviewed by
Jason Pirozzolo, DO
Fellowship Trained and Board Certified in Sports Medicine

Last medically reviewed: November 1, 2026